




Bulkamid™
Urethral Bulking System

Ready to say
goodbye to
bladder leaks?

Say hello to **Bulkamid**

Bulkamid is a minimally invasive
and long-lasting treatment option
for bladder leaks caused by stress
urinary incontinence (SUI).



Do you struggle with bladder leaks when you...



Sneeze



Exercise



Laugh




Cough



Walk

You're
not alone



~1 in 4 women
in europe experience
urinary incontinence.⁴

What is stress urinary incontinence (SUI)?

Stress urinary incontinence (SUI), some-times called stress incontinence, is a condition where unwanted bladder leaks happen during normal daily activities such as coughing, laughing, or exercise.

How can Bulkamid help?

Bulkamid is a urethral bulking agent that is used to treat stress urinary incontinence. It is a soft, water-based gel that can be used to restore the natural closing of the urethra. Similar to facial fillers, Bulkamid adds volume to the surrounding tissue.¹

Discover the benefits of Bulkamid

► Safe and effective

Bulkamid is a safe and effective treatment that can be performed in an office setting or as an outpatient procedure.

► The procedure

The procedure is a series of 3 – 4 injections that takes about 20 minutes under local anesthetic.²

► Long-lasting relief

Bulkamid is designed to effectively treat SUI in women, with clinical support out to seven years.³

In one study,
64% of women
chose Bulkamid
to treat their SUI
over other options.²



Don't settle for frustrating urinary symptoms or incontinence pads.

Ask your doctor today if Bulkamid is right for you.

To learn more about SUI visit: www.urologymatters.com

1. Chapple C, Dmochowski R. Particulate versus non-particulate bulking agents in the treatment of stress urinary incontinence. Res Rep Urol. 2019; 11: 299–310.
2. Dwyer L, Weaver E, Rajai A, Cox S, Reid F. "Voice Your Choice": A study of women's choice of surgery for primary stress urinary incontinence. Int Urogynecol J. 2020;31:769–777.
3. Brosche T, Kuhn A, Lobodasch K, Sokol ER. Seven-year efficacy and safety outcomes of Bulkamid for the treatment of stress urinary incontinence. Neurourol Urodyn. 2021;40:502–508.
4. European Association of Urology. (2024). The health, socio-economic and environmental costs of continence problems in the EU.
https://d56bochluxqnz.cloudfront.net/media/Socio-economic_report_UrgetoAct.pdf

Important Safety Information

Bulkamid Urethral Bulking System is intended to be used as a urethral bulking agent for the treatment of female urinary incontinence, or the involuntary loss of urine with movement or activity that puts pressure on the bladder, such as coughing or sneezing. Patients who have an infection of the bladder or urethra, active herpes outbreak, pelvic infection, or a history of damage to the urethral tissue may not be a candidate for a Bulkamid procedure.

With all medical procedures, there are risks associated with the procedure and the use of the device, including long-term implantation of the hydrogel. These risks include but are not limited to pain with urination, blood in the urine, urinary tract infection, or difficulty urinating immediately following the procedure. Long-term risks include difficulty urinating, abscess, scarring of the urethra, and urinary urgency.

Be sure to talk with your doctor so that you thoroughly understand all of the risks and benefits associated with the use of the device.

Results from case studies are not necessarily predictive of results in other cases. Results in other cases may vary.

CAUTION: The law restricts these devices to sale by or on the order of a physician. Indications, contraindications, warnings and instructions for use can be found in the product labelling supplied with each device. Products shown for INFORMATION purposes only and may not be approved or for sale in certain countries. This material not intended for use in France.

**Boston
Scientific**
Advancing science for life™

www.bostonscientific.eu

© 2026 Boston Scientific Corporation or its affiliates. All rights reserved.

URO-2237808-AB
CE 0123